

Short note on the Rabi model

K. Ziegler
*Institut für Physik, Universität Augsburg,
 D-86135 Augsburg, Germany*
 and
*Physics Department,
 New York City College of Technology,
 CUNY, Brooklyn, NY 11201, USA*
 (Dated: March 2, 2013)

The spectral density of the Rabi model is calculated exactly within a continued fraction approach. It is shown that the method yields a convergent solution.

PACS numbers: 03.65.Ge, 02.30.Ik, 42.50.Pq

A recent work by Braak [1] has renewed the interest in the old problem of coupling a photon field to a single spin 1/2 state, using the Rabi model [2]. The central statement of this work is that the eigenfunctions in Bargmann representation must be analytic functions in the entire complex plane. Based on this condition, a procedure is derived from the series expansion of the eigenstates which provides a recursive evaluation of the spectrum. It is claimed that the series expansion yields an exact solution of the Rabi model, which cannot be obtained from a direct continued fraction (CF) approach. In the following it is shown that this statement is not valid, and that the use of the extra condition of analyticity of the eigenfunction in Bargmann representation is not necessary. Therefore, the CF is directly applicable to the Rabi model and yields an exact solution for the spectral density, where the term “exact” means that the CF is convergent and the spectral density can be evaluated by a simple algorithm to any desired accuracy.

The Rabi model is defined by the Hamiltonian

$$H_R = \omega a^\dagger a + \Delta \sigma_3 + U(a^\dagger + a)\sigma_1, \quad (1)$$

where $a^\dagger$ (a) are creation (annihilation) operators of a photon, the Pauli matrices σ_j ($j = 0, \dots, 3$) describe operations on the spin-1/2 state, and Δ is a symmetry breaking field for the spin. U is the coupling strength between the photons and the spin states. This Hamiltonian maps a product state $|N\rangle \otimes |\sigma\rangle$ with N photons to $|N \pm 1\rangle \otimes |\sigma'\rangle$, where $\sigma, \sigma' = \uparrow, \downarrow$. Thus eigenstates of H are superpositions of product states, which can be derived by a recursive approach for the coefficients of the superposition [3]. In the following we apply the recursive projection method (RPM) of Ref. [4]. This method is based on a random walk expansion in the underlying Hilbert space which visits each subspace at most once [5]. Consequently, there are no loops in the random walk, which leads directly to a CF.

The RPM, as described in Ref. [4], can be directly applied to the resolvent $\langle N; s | (z - H)^{-1} | N; s' \rangle$ and yields immediately the spectral density with respect to $|N; s\rangle$:

$$\rho_N(E) = \frac{1}{2} \sum_{s=\uparrow, \downarrow} \lim_{\epsilon \rightarrow 0} \text{Im}(\langle N; s | (E - i\epsilon - H)^{-1} | N; s \rangle). \quad (2)$$

The simplest case is the resolvent $\langle 0; s | (z - H)^{-1} | 0; s' \rangle$ of states without photons $|0, s\rangle$ and with spin projection $s = \uparrow, \downarrow$:

$$(\langle 0; s | (z - H)^{-1} | 0; s' \rangle) = \lim_{n \rightarrow \infty} \begin{pmatrix} g_n & 0 \\ 0 & h_n \end{pmatrix}, \quad (3)$$

where the eigenvalues of H are the poles of the resolvent. Then the matrix elements g_n, h_n are subject to the following recurrence relations

$$g_k = \frac{1}{z - \omega(n - k) + \Delta - U^2(n - k + 1)h_{k-1}} \quad (4)$$

$$h_k = \frac{1}{z - \omega(n - k) - \Delta - U^2(n - k + 1)g_{k-1}} \quad (5)$$

for $1 \leq k \leq n$ with the initial values

$$g_0 = \frac{1}{z - \omega n - \Delta}, \quad h_0 = \frac{1}{z - \omega n + \Delta}. \quad (6)$$

The iteration of these recurrence relations gives a finite CF of the standard form [6]

$$g_n = b_0 + \frac{a_1}{b_1 + \frac{a_2}{b_2 + \dots}} \equiv b_0 + \frac{a_1}{|b_1|} + \dots + \frac{a_n}{|b_n|} \quad (7)$$

with coefficients

$$a_1 = 1, \quad a_k = (1 - k)U^2, \quad b_0 = 0, \quad b_1 = z - \Delta, \quad b_k = z - (k - 1)\omega + (-1)^k \Delta \quad (2 \leq k \leq n). \quad (8)$$

For the corresponding coefficients of h_n we must only replace Δ by $-\Delta$. Therefore, all considerations for g_n apply to h_n after replacing $\Delta \rightarrow -\Delta$. Due to $b_0 = 0$ the inverse of g_n can be obtained directly by inverting g_n in Eq. (7):

$$a_1/g_n = b_1 + \frac{a_2}{|b_2|} + \dots + \frac{a_n}{|b_n|}. \quad (9)$$

This means that the poles of g_n are obtained from the zeros of the related CF, and vice versa.

By taking the limit $g = \lim_{n \rightarrow \infty} g_n$ the CF in Eq. (7) or (9) yields the spectral density of the Rabi model for the complex energy z . This requires the limit $n \rightarrow \infty$ though, whose existence is a consequence of the Pringsheim Theorem (cf. [6]): Considering the tail t_n of the CF

$$g = b_0 + \frac{a_1}{|b_1|} + \dots + \frac{a_n}{|b_n|} + t_n \quad (10)$$

with

$$t_n = \frac{a_{n+1}}{|b_{n+1}|} + \dots, \quad (11)$$

we estimate the error when we truncate g at $k = n$ (which gives g_n). The Pringsheim Theorem states that for complex a_k, b_k with $|b_k| \geq |a_k| + 1$ ($k \geq n$) the truncated CF $t_{n,l}$

$$t_{n,l} = \frac{a_{n+1}}{|b_{n+1}|} + \dots + \frac{a_l}{|b_l|} \quad (n + 1 < l) \quad (12)$$

is convergent for $l \rightarrow \infty$ and converges to a t_n with

$$0 \leq |t_n| \leq 1. \quad (13)$$

For the coefficients of the Rabi model in Eq. (8) the conditions for the coefficients are satisfied for a given z and for sufficiently large n if $U^2 < \omega$. Then (13) means that the ratio t_n/b_n is $O(1/n)$, which implies

$$t_{n-1} = \frac{a_n}{b_n + t_n} = \frac{a_n}{b_n} + O(1/n) = \frac{(1 - n)U^2}{z - (n - 1)\omega - (-1)^n \Delta} + O(1/n) = \frac{U^2}{\omega} + O(1/n). \quad (14)$$

This result enables us to approach the exact solution with a sequence of truncated CF's with a given error $O(1/n)$:

$$g = b_0 + \frac{a_1}{|b_1|} + \dots + \frac{a_{n-1}}{|b_{n-1}|} + \frac{a_n/b_n + O(1/n)}{|1|}. \quad (15)$$

The finite CF can also be expressed as the ratio

$$g = \frac{A_{n-1} + [a_n/b_n + O(1/n)]A_{n-2}}{B_{n-1} + [a_n/b_n + O(1/n)]B_{n-2}}, \quad (16)$$




FIG. 1: Spectral density $\rho_0(E)$ for maximally $n = 500$ photons and $\Delta = 0.4$, $U = 0.99$, $\epsilon = 0.005$ in units of the photon frequency ω . The green (red) curves are contributions from g_n (h_n).

where A_{n-l} (B_{n-l}) are polynomials in z of order $n-l-1$ ($n-l$) which are generated by the recurrence relations (cf. [6])

$$A_k = b_k A_{k-1} + a_k A_{k-2}, \quad B_k = b_k B_{k-1} + a_k B_{k-2} \quad (k \geq 1) \quad (17)$$

with initial conditions $A_{-1} = 1$, $A_0 = b_0 = 0$, $B_{-1} = 0$, $B_0 = 1$.

Conclusions and remarks: Eq. (16) allows us to approach successively the spectral density of the Rabi Hamiltonian, where the accuracy is improved with increasing n . The error is estimated by the tail t_n of (14). According to our definition this provides an exact solution of the Rabi spectrum, where the poles of g and h are the corresponding eigenvalues. The evaluation of $\rho_N(E)$ for $N > 0$ can be performed within the same approach.

The individual matrix elements g , h avoid level crossing due to parity conservation, since eigenstates of consecutive eigenvalues have different parity. However, the sum of g_n and h_n in the spectral density $\rho_0(E)$ provides level crossing by shifting their levels relative to each other, for instance, by changing Δ .

The “rotating-wave” approximation of the Rabi Hamiltonian yields the Jaynes-Cummings Hamiltonian

$$H_{JC} = \omega a^\dagger a + \Delta \sigma_3 + U(a^\dagger \sigma_- + a \sigma_+) \quad (18)$$

with $\sigma_\pm = (\sigma_1 \pm i\sigma_2)/2$. This Hamiltonian does not belong to the class of models with recursive solutions because the recurrence relation of the RPM terminates already after the first step for each k :

$$g_k = \frac{z - \omega(n-k+1) - \Delta}{[z - \omega(n-k+1)]^2 - \Delta^2 + \omega - U^2(n-k+1)}$$

$$h_k = \frac{1}{z - \omega(n-k) - \Delta}. \quad (19)$$

Here the evaluation of the eigenvalues requires only the solution of a quadratic equation for each k , in agreement with the original work on this model [7, 8].

Acknowledgments

The author is grateful for the hospitality at the New York City College of Technology during his sabbatical.

-
- [1] D. Braak, Phys. Rev. Lett. **107**, 100401 (2011).
 - [2] I.I. Rabi, Phys. Rev. **49**, 324 (1936); **51**, 652 (1937).
 - [3] S. Schweber, Ann. Phys. (N.Y.) **41**, 205 (1967).
 - [4] K. Ziegler, J. Phys. B: At. Mol. Opt. Phys. **44**, 145302 (2011).
 - [5] K. Ziegler, Phys. Rev. A **81**, 034701 (2010).
 - [6] O. Perron, *Die Lehre von den Kettenbrüchen*, Teubner Verlag (Leipzig, Berlin 1912).
 - [7] E.T. Jaynes and F.W. Cummings, Proc. Inst. Elect. Eng. **51**, 89 (1963).
 - [8] F.W. Cummings, Phys. Rev. **140**, A1051 (1965).